

PATENT ABSTRACTS OF JAPAN

(11) Publication number : 11-279159
 (43) Date of publication of application : 12. 10. 1999

(51) Int. Cl.

C07D241/04
 C07B 57/00
 // C07M 7:00

(21) Application number : 10-095385
 (22) Date of filing : 24. 03. 1998


(71) Applicant : NIPPON SODA CO LTD
 (72) Inventor : TAKADA MITSUMASA
 KATO TETSUYA
 HIRAI YUKIO

(54) PRODUCTION OF OPTICALLY ACTIVE PIPERAZINECARBOXYLIC ACID ESTER


(57) Abstract:

PROBLEM TO BE SOLVED: To produce the subject compound in high yield by heating the S-isomer or a solution consisting essentially of the S-isomer in the presence of a base, providing a mixture of the R- and the S-isomers and then optically resolving the resultant mixture.


SOLUTION: L-Menthyl (S)-1,4-bis(phenylmethyl)-2-piperazinecarboxylate represented by formula I or a solution consisting essentially of the menthyl ester is heated in the presence of a base to provide a mixture of L-menthyl (R)-1,4-bis(phenylmethyl)-2-piperazinecarboxylate represented by formula II with the L-menthyl (S)-1,4-bis(phenylmethyl)-2-piperazinecarboxylate. The resultant mixture is then optically resolved to thereby produce L-menthyl (R)-1,4-bis(phenylmethyl)-2-piperazinecarboxylate represented by formula III. The compound represented by formula III is useful as an intermediate for (R)-4-(3-phosphono-2-propenyl)-2-piperazinecarboxylic acid represented by formula IV which is an excellent antidepressant and anti-Alzheimer disease agent.




I



II



III



IV

LEGAL STATUS

[Date of request for examination]

[Date of sending the examiner's decision of rejection]

[Kind of final disposal of application other than the examiner's decision of rejection or application converted registration]

[Date of final disposal for application]

[Patent number]

[Date of registration]

[Number of appeal against examiner's decision of rejection]

[Date of requesting appeal against examiner's decision of rejection]

The Japanese Patent Office is not responsible for any damages caused by the use of this translation.

1. This document has been translated by computer. So the translation may not reflect the original precisely.
2. **** shows the word which can not be translated.
3. In the drawings, any words are not translated.

DETAILED DESCRIPTION


[Detailed Description of the Invention]

[0001]

[The technical field to which invention belongs] this invention relates to the useful manufacture technique industrially [(R)-1 of the optical activity shown by the formula (III), and 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester]. The (R)-4-(3-phosphono-2-propenyl)-2-piperazine carboxylic acid shown by the formula (V) is outstanding anti-**** and anti-Alzheimer's-disease medicine (JP,63-203691,A) which were developed at the Switzerland sand company, and the compound expressed with a formula (III) is useful as intermediate field of (V).


[0002]

[Formula 5]



[0003]

[Formula 6]




[0004]

[Description of the Prior Art] Conventionally, after the synthesis method of (R)-1 and 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester (R bodies) expressed with a formula (III) made the transesterification of 1 and 4-screw (phenyl methyl)-2-piperazine carboxylic-acid low-grade alkyl ester and L-menthol perform, according to the solubility difference of the obtained diastereomer compound, it divided the diastereomer and had obtained R bodies (III). At that time, (S)-1 expressed with the formula (I) which is the principal component of a mother liquor, and 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester (S bodies) became unnecessary, and yield was also low (JP,63-203691,A).

[0005]

[Formula 7]



[0006]

[Problem(s) to be Solved by the Invention] The development of the technique of reusing S bodies (I) which became unnecessary from the above thing was desired. this invention carries out racemization of the S unnecessary bodies (I), and is made into R and S mixture, and R bodies (III) are further taken out from the mixture, and it aims at measuring the enhancement in yield.


[0007]

[Means for Solving the Problem] This invention persons came to complete this invention, as a result of examining the above-mentioned technical problem.

[0008] That is, it is the manufacture technique of R bodies (III) characterized by carrying out optical resolution of it after this invention heats the solution which makes S bodies or S bodies a principal component under base presence and makes it R and S mixture (II).

[0009]

[Formula 8]



[0010] The bases under the base presence of this invention are an organic base and an inorganic base, and are inorganic bases preferably, for example, a sodium hydride, a sodium methylate, and a t-butoxy potassium are mentioned.

[0011] To S bodies (I), although it is a grade mol theoretically, a reaction may advance also in the amount of catalysts, for example, 10 - 20 mol% is sufficient as the amount of the base used of this invention to (I).

[0012] A solvent inactive for a reaction about the solvent of this invention is desirable, for example, are benzene and toluene.

[0013] Temperature was applied about the reaction temperature of this invention, and it is [direction] good and reacts at 110-120 degrees C preferably.

[0014] Although it goes on about the reaction time of this invention for a short time, it reacts preferably for 2 to 3 hours.

[0015] It repeats several times that the crystallization conditions of this invention add and carry out the crystallization of the organic solvent to the hydrochloric-acid aqueous solution. By this, R (III) with more sufficient purity can be obtained.

[0016] The organic solvent of the crystallization conditions of this invention is a polar solvent, for example, are the ether and ethanol.

[0017] Racemization of the S bodies (I) of this invention can be carried out, and the technique of obtaining R bodies (III) can be recycled any number of times.

[0018] In order to perform the manufacture technique of this invention industrially, the following two technique is possible.

[0019] (1) Perform optical resolution after carrying out racemization, making transesterification perform in addition to the reaction mixture of N [which is expressed with a formula (IV) in the solution which makes a principal component S bodies (I) or S bodies (I)], and N'-dibenzyl piperazine-2-carboxylic-acid low-grade alkyl ester, and L-menthol.

[0020] (2) In addition to the diastereomer compound solution which the compound and L-menthol which are expressed with a formula (IV) in this solution are made to react, and was obtained, perform optical resolution simultaneously after heating and carrying out racemization of the S bodies (I).

[0021] It is recyclable any number of times as it stated the account of a front, when which technique was adopted above.

[0022] An example is given to below and this invention is explained to it in detail.

[0023]

[Example] (R) -1, synthetic :N [of 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester], and N'-dibenzyl piperazine-2-carboxylic-acid methyl-ester 36.7g (114mmol) was agitated, using toluene 100ml as a solvent, L-menthol 26.1g (167mmol) was added at the room temperature, heating reflux was performed for 30 minutes using the Dean Stark trap, and toluene was dehydrated. NaH(distributed object, 55 - 60%)1g (26mmol) was added after cooling a reaction mixture, and heating reflux was performed for 3 hours. At this time, 60ml toluene was distilled off and the ** methanol was performed by the technique of adding new toluene 60ml. 2N after cooling a reaction mixture HCl When 60ml and ether 150ml were added and churning was carried out at 20-25 degrees C for 1 hour, the crystal separated, and (R)-1 and 32.4g (diastereomer purity 65.8% d.) of 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester-hydrochloride-hydrates were obtained by ****ing the crystal and drying.

[0024] Furthermore, it is 0.2N about this mixture. HCl 60ml, EtOH Recrystallization was performed by 100ml and, finally (R)-1 and 20.2g (35% [of yield] and diastereomer purity 98.8%d.e.) of 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester-hydrochloride-hydrates were obtained.

[0025] At this time, NaOH was extracted 3 times by toluene 30ml as optimum-dose **** and pH 12 28% to the filtrate of two obtained batches. Concentration hardening by drying of the organic layer was dried and carried out with magnesium sulfate, and it obtained oil 38.7g (R/S=2/7). Toluene 100ml was added to this oil, heating reflux was performed for 30 minutes using D an Stark, and toluene was dehydrated. NaH(distributed object, 55 - 60%)1g (26mmol) was added after cooling a reaction mixture, and heating reflux was performed for 3 hours. (When the reaction mixture was checked at this time, racemization was carried out to (R/S=1 / 1).) 2N By adding HCl60ml and ether 150ml, a crystal's separating, if it agitates at 20-25 degrees C for 1 hour, and ****ing the crystal, and drying (R) -1 and 17.6g (diastereomer purity 75.0%d.e.) of 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester-hydrochloride-hydrates were obtained.

[0026] Furthermore, it is 0.2N about this mixture. HCl 60ml, EtOH Recrystallization was performed by 100ml and, finally (R)-1 and 12.0g (21% [of yield] and diastereomer purity 96.1%d.e.) of 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester-hydrochloride-hydrates were obtained. Consequently, the total yield which added one racemization recycling improved with Y= 56.2% (32.2g).

[0027]

[Effect of the Invention] The manufacturing methods of this invention are optical activity (R)-1 [useful as intermediate field of the drug of outstanding anti-**** and anti-Alzheimer's-disease medicine (JP,7-23387,A)], and the industrially excellent manufacture technique which can manufacture efficiently 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester.

[Translation done.]

NOTICES

The Japanese Patent Office is not responsible for any damages caused by the use of this translation.


1. This document has been translated by computer. So the translation may not reflect the original precisely.
2. **** shows the word which can not be translated.
3. In the drawings, any words are not translated.

CLAIMS


[Claim(s)]

[Claim 1] The solution which makes a principal component (S)-1 expressed with a formula (I), 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester, or this ester is heated under base presence. After considering as (R)- expressed with a formula (II), (S)-1, and 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester mixture, (R)-1, the manufacture technique of 4-screw (phenyl methyl)-2-piperazine carboxylic-acid-L-menthyl ester which are expressed with the formula (III) characterized by carrying out optical resolution.


[Formula 1]



[Formula 2]



[Formula 3]




[Claim 2] The manufacture technique according to claim 1 of considering as the compound expressed with a formula (II), performing transesterification in addition to the reaction mixture of N [which is expressed with a formula (IV) in the solution which makes a principal component the compound or this compound expressed with a formula (I)], and N'-dibenzyl piperazine-2-carboxylic-acid low-grade alkyl ester, and L-menthol, and performing a chemistry split.

[Claim 3] The manufacture technique according to claim 1 which carries out optical resolution simultaneously in addition to the reaction mixture to which the transesterification of the compound and L-menthol which are expressed with a formula (IV) was made to perform after considering as the compound which heats the solution which makes a principal component the compound or this compound expressed with a formula (I) under base presence, and is expressed with a formula (II).

http://www4.ipdl.jpo.go.jp/cgi-bin/tran_web_cgi_ejje?u=http%3A%2F%2Fwww.ipdl.jpo... 2001-08-21

[Formula 4]



(R shows a low-grade alkyl group among a formula.)

[Translation done.]